

Intertrigo is an infection caused by a yeast condition called *Candida*. It comes from the same family of yeast infections that cause Athletes Foot, Jock Itch and Nappy Rash

Intertrigo is very common. This leaflet is designed to raise awareness of the condition and to aid self-help treatment.

For further information or advice contact your local doctor, pharmacy or breast screening unit.

#### Contact Information

**King's Lynn Breast Screening Unit**  
**01553 613662**  
(Open 8.30am-4.30pm Mon - Fri)

**Leaflet researched and designed by:-**  
Marilyn O'Connell (AP Breast Screening)  
The West Norfolk Breast Care Unit,  
The Queen Elizabeth Hospital,  
King's Lynn NHS Foundation Trust,  
Norfolk PE30 4ET

Thanks to the Queen Elizabeth Hospital Breast Screening Team and the King's Lynn Breast Screening ladies for their help and support in the completion of this leaflet.

# Understanding Under-breast Soreness (Intertrigo)



## Symptoms and Treatment

## What causes Under-breast Soreness

Under-breast soreness usually occurs in women who are larger breasted or very sporty.

The main causes are sweat rash, 'skin to skin' rubbing and the growth of yeast, which appear as a rash in the skin fold, under the breast. This is commonly known as Intertrigo.

If untreated, Intertrigo can spread to other parts of the body or worsen. In some cases it can lead to creamy coloured discharge and emit a pungent smell.

## What are the symptoms of Intertrigo?

The main symptoms are:

- Inflammation
- Soreness / Discomfort
- Split skin
- Spots
- Itching
- Discharge (often pungent)



## What is the most effective treatment?

The most effective treatment for Under-breast Soreness is to:

- Wash the affected area morning and night and pat dry, rather than rub dry. Wear good supportive bras (cotton is known to be effective). This will help stop the 'skin to skin' rubbing. (Nylon bras are known to make the condition worse)
- Do not share flannels or towels, to avoid the spread of infection.
- In cases where the skin is very red, very inflamed and sore, ask your pharmacist for a \*Hydrocortisone Cream.
- In cases where the skin is red, sore or itchy, ask your pharmacist for an \*Anti-fungal cream or powder.
- Once the infection has cleared, it may be advisable to use a \*Barrier cream (i.e. zinc and castor oil) on a regular basis to help stop the infection returning.
- If none of the above treatments help or your symptoms worsen, make an appointment to see your doctor.

\* Hydrocortisone, anti-fungal creams/powders and barrier creams are all available over the counter, please follow instructions.